

land

ington,

1848 from
nd (Owen,
f elements
the species

und in the
sing a new
o those of
nen which
e from the
es (Owen,
Greenway,


y of London

erreaux and des Murs) are known from New
 etids are considered to be allied to the rails and
 land. *R. orarius* was described from Quaternary
 by comparison of dimensions to be larger than
 1989).

ods

erial

has been collected from swamp and cave sites across
 1984; Beauchamp & Worthy, 1988) (Fig. 1). Bones
 o between 25,000 and 1000 years bp (Yaldwyn, 1958;



the Quaternary fossil locations, and (●) sites where material

Horn, 1967
 provided
 from re-
 populati-
 the Sout-

Under
 elements
 graduate
 Papa Te-
 excludin-
 latero-m-
 Worthy,
 Sterna: p-
 above th-
 breadth
 tip (Prer-
 (Pelvis I-
 width ac-
 posterio-

Data
 rare with
 collected
 be made

Data v-
 each of t-
 of the t-
 measure-
 was proc-
 further a-
 (Bartlett'
 (Fig. 5b)
 SI) 1—1
 Marlbor-
 should be
 several si-
 pukeko
 compare-

Data f-
 ratios of
 and com-

Mean

and
reflect
y at
from
are

the
p. D.
of
tern
are

ame
the
rity
e SI
unt



Tarsometatarsus DB.

Mean
n the

22

St

—

Bo

Pr

C.

C.

H

H

H

H

Ci

St

Pe

Pe

Pe

Pe

Fe

Fe

Fe

Fe

Ti


Ti

Ti

Ti

Ta

Ta



The n
terminal
proporti
metatars
despite t
emphasi
The inter
takahe.

The at
birds is
(Table I
significai
respectiv

Tarsor
Bartlett's
indicatin
differenc
Within th
scores in
differed v
NI and S
expected

5
5
5
5


0
0
0
0

ling to

nd SI
n the
otted
22:36,
2:496,
cture.
ng in
ate a
very
its of
I and
ongly
iance
that
m SI

(Inter
 Overal
 from t
 Exam
 reveale
 than w
 of the
 others
 larger
 25% b
 such th

FIG. 6
 SI (■) ■
 $r^2 = 0.006$
 0.3256*
 22.7846,
 Lntib =



2.5



2.45

2:

L
%

—
B
—
F
T
T
—

b
o
b
m
d
d
v

nc

%

val
'sm

certain
they can
emphen
Pacific
ization.
ne wide
Zealand
cases,
stdates

, 1979;
lt of an
erefore,
o or an
whether
ightless

warrant
, it has
nergetic
ata for
ightless
to two
icult to
ant and
n to fix
olocene
ological
taking
ns and
nabitat.
in par-
a wide
be the
orms in
nometical
state is
as been
y might

esented
onizing
f flight-
bserved
takahe.
closely

with the principles of the 0-666 power of power functions. E (legs for instance) between phylogenetic components and the of this relationship

The two forms of kagu are very similar scaled versions of o diverged from an a larger size. In contrast must have been in r forms across the r crossed the Cook specializing selective

ANCOVA analysis form than either is SI takahe, the NI ta and generally would suggest that the two takahe may represent most from the pul therefore imply the large size and a flight

Examination of the New Zealand may from fossil bones (slighter) but not quite another large blue leaving no physical (1962) observed a v remain. This 'White' being "easily struck" inability of the species ancestors have given what may transpire similar.

The application of In the meantime, the the original names *Porphyrio mantelli*

of constraints such as requirements of maintenance in different ancestor in different species would seem more closely related to similar.

For their comments on the paper at the Waitomo Caves and Te Papa Tongarewa; Canterbury Museum.

183: 125–146.

Lawler, D. M. (Ed.). New York:

Malden. *Smithson. Contribs*

Myiophyrio (= *Notornis*) *mantelli*:

Contributions. *Trans. Proc. R. Soc.*

London: Chapman and University Press.

1904. *Trans. Proc. N.Z. Inst.* **24**:

10.

in New Zealand. *Notornis* **27**:

1904. *Notornis* of *Oxyura australis* (blue-

1901–394.

London: Reed.

scour and aligning basins at a

1–38.

London: Chapman and University Press.


Nat. **109**: 547–563.

Nat. **144**: 628–642.

University of Auckland, New

Myzomela, a new genus of wren

(Waitomo) cave, Waitomo. *J. R. Soc.*



1;
2;
3;
4;
5;
6;
7;
8;
9;
10;

11;
12;
13;